

## LA-UR-21-21790

Approved for public release; distribution is unlimited.

Title: Variational calculation of closed-shell nuclei up to  $A=40$

Author(s): Lonardonì, Diego

Intended for: One-Page Publication Highlights

Issued: 2021-02-23

---

**Disclaimer:**

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by Triad National Security, LLC for the National Nuclear Security Administration of U.S. Department of Energy under contract 89233218CNA000001. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.


# Variational calculation of closed-shell nuclei up to $A = 40$

## Objectives


- Compute ground state properties of closed-shell nuclei up to  $A = 40$  with quantum Monte Carlo.
- Analyze the behavior of phenomenological Hamiltonians in medium-heavy nuclei.
- Study the high-momentum components of the nuclear wave function and potential in-medium modifications of the nucleon form factors.

## Impact

- Observe a change in the behavior of the UIX 3-body force — from attractive to repulsive — already in relatively small nuclear systems, such as  $^{16}\text{O}$ .
- Confirm the universality of the tail of the momentum distribution for a given class of Hamiltonians.
- Show very little  $A$  dependence of the Coulomb sum rule for  $A \geq 12$ : no evidence of in-medium modification of the nucleon form factor.



Longitudinal elastic form factor in  $^{16}\text{O}$  for different phenomenological Hamiltonians. The UIX 3-body force combined with the AV18 potential provides a good description of the structure of the nucleus.



Coulomb sum rules for  $A \leq 40$ . Symbols with error bars show GFMC calculations employing the AV18+IL7 potential. The curves show CVMC results for AV18+UIX.

## Accomplishments

- Development of a cluster variational Monte Carlo approach (CVMC) to study the properties of medium-heavy nuclei employing realistic nuclear interactions.
- Analysis of binding energies, charge radii, one- and two-body densities, one-body momentum distributions, charge form factors, and Coulomb sum rules for  $A = 4, 16, 40$ .



U.S. DEPARTMENT OF  
**ENERGY**

Office of  
Science

**NUCLEI**  
Nuclear Computational Low-Energy Initiative

**References:** [Phys. Rev. C 96, 024326 \(2017\)](#)

**Contact:** D. Lonardoni  
[lonardoni@nscl.msu.edu](mailto:lonardoni@nscl.msu.edu)